

**Sanofi Pasteur**  
**306 – Adacel®**

**Full Prescribing Information**

**HIGHLIGHTS OF PRESCRIBING INFORMATION**

These highlights do not include all the information needed to use Adacel safely and effectively. See full prescribing information for Adacel.

**Adacel (Tetanus Toxoid, Reduced Diphtheria Toxoid and Acellular Pertussis Vaccine Adsorbed)**

**Suspension for Intramuscular Injection**

**Initial US Approval: 2005**

**RECENT MAJOR CHANGES**

Warnings and Precautions. (5.2) 09/2017

**INDICATIONS AND USAGE**

Adacel is a vaccine indicated for active booster immunization against tetanus, diphtheria and pertussis. Adacel is approved for use as a single dose in persons 10 through 64 years of age. (1)

**DOSAGE AND ADMINISTRATION**

A single intramuscular injection of 0.5 mL. (2.1)

**DOSAGE FORMS AND STRENGTHS**

Single-dose vials and prefilled syringes containing a 0.5 mL suspension for injection. (3)

**CONTRAINDICATIONS**

Severe allergic reaction (eg, anaphylaxis) to any component of Adacel or any other diphtheria toxoid, tetanus toxoid and pertussis antigen-containing vaccine. (4.1)

Encephalopathy (eg, coma, decreased level of consciousness, prolonged seizures) within 7 days of administration of a previous pertussis antigen-containing vaccine. (4.2)

**WARNINGS AND PRECAUTIONS**

For one presentation of Adacel, the tip caps of the prefilled syringes may contain natural rubber latex, which may cause allergic reactions in latex sensitive individuals. (5.2, 16)

If Guillain-Barré syndrome occurred within 6 weeks of receipt of a prior vaccine containing tetanus toxoid, the risk for Guillain-Barré syndrome may be increased following a subsequent dose of Adacel vaccine. (5.3)

Progressive or unstable neurologic conditions are reasons to defer Adacel vaccination. (5.4)

Persons who experienced an Arthus-type hypersensitivity reaction following a prior dose of a tetanus toxoid-containing vaccine should not receive Adacel unless at least 10 years have elapsed since the last dose of a tetanus toxoid-containing vaccine. (5.5)

Syncope (fainting) can occur in association with administration of injectable vaccines, including Adacel. Procedures should be in place to prevent falling injury and manage syncopal reactions. (5.7)

**ADVERSE REACTIONS**

The most common solicited injection site reactions occurring within 0-14 days following vaccination with Adacel were:

- For Adolescents 11-17 years of age: pain (77.8%), swelling (20.9%), erythema (20.8%).

- For Adults 18-64 years of age: pain (65.7%), swelling (21.0%), erythema (24.7%). (6.1)

The most common solicited systemic reactions occurring within 0-14 days following vaccination with Adacel were:

- For Adolescents 11-17 years of age: headache (43.7%), body ache or muscle weakness (30.4%), tiredness (15.1%).

- For Adults 18-64 years of age: headache (33.9%), body ache or muscle weakness (21.9%). (6.1)

**To report SUSPECTED ADVERSE REACTIONS, contact Pharmacovigilance Department, Sanofi Pasteur Inc., Discovery Drive, Swiftwater, PA 18370 at 1-800-822-2463 (1-800-VACCINE) or VAERS at 1-800-822-7967 or <http://vaers.hhs.gov>.**

**DRUG INTERACTIONS**

When Adacel vaccine was administered concomitantly with trivalent inactivated influenza vaccine (TIV) to adults 19-64 years of age, a lower antibody response was observed for pertactin antigen as compared to Adacel vaccine administered alone. (7.1, 14.3)

Immunosuppressive therapies may reduce the immune response to Adacel. (7.2)

Do not mix Adacel vaccine with any other vaccine in the same syringe or vial.

**USE IN SPECIFIC POPULATIONS**

Safety and effectiveness of Adacel vaccine have not been established in pregnant women. (8.1)

Pregnancy Surveillance Registry: contact Sanofi Pasteur Inc. at 1-800-822-2463 (1-800-VACCINE). (8.1)

See 17 PATIENT COUNSELING INFORMATION

Revised: [XXX/2017]

**FULL PRESCRIBING INFORMATION: CONTENTS\***

**1 INDICATIONS AND USAGE**

**2 DOSAGE AND ADMINISTRATION**

- 2.1 Preparation for Administration
- 2.2 Administration, Dose and Schedule
- 2.3 Additional Dosing Information

**3 DOSAGE FORMS AND STRENGTHS**

**4 CONTRAINDICATIONS**

- 4.1 Hypersensitivity
- 4.2 Encephalopathy

**5 WARNINGS AND PRECAUTIONS**

- 5.1 Management of Acute Allergic Reactions
- 5.2 Latex
- 5.3 Guillain-Barré Syndrome and Brachial Neuritis
- 5.4 Progressive or Unstable Neurologic Disorders
- 5.5 Arthus-Type Hypersensitivity
- 5.6 Altered Immunocompetence
- 5.7 Syncope

**6 ADVERSE REACTIONS**

- 6.1 Clinical Trials Experience
- 6.2 Postmarketing Experience

**7 DRUG INTERACTIONS**

- 7.1 Concomitant Vaccine Administration
- 7.2 Immunosuppressive Treatments

**8 USE IN SPECIFIC POPULATIONS**

- 8.1 Pregnancy

8.3 Nursing Mothers

8.4 Pediatric Use

8.5 Geriatric Use

**11 DESCRIPTION**

**12 CLINICAL PHARMACOLOGY**

- 12.1 Mechanism of Action

**13 NON-CLINICAL TOXICOLOGY**

- 13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

**14 CLINICAL STUDIES**

- 14.1 Immunological Evaluation in Adolescents and Adults, 10 Through 64 Years of Age
- 14.2 Concomitant Hepatitis B Vaccine Administration
- 14.3 Concomitant Influenza Vaccine Administration

**15 REFERENCES**

**16 HOW SUPPLIED/STORAGE AND HANDLING**

**17 PATIENT COUNSELING INFORMATION**

\* Sections or subsections omitted from the full prescribing information are not listed.

1 **FULL PRESCRIBING INFORMATION:**

2 **1 INDICATIONS AND USAGE**

3 Adacel® is a vaccine indicated for active booster immunization against tetanus, diphtheria and  
4 pertussis. Adacel vaccine is approved for use as a single dose in individuals 10 through 64 years  
5 of age.

6 **2 DOSAGE AND ADMINISTRATION**

7 **2.1 Preparation for Administration**

8 Just before use, shake the vial or syringe well until a uniform, white, cloudy suspension results.

9 Parenteral drug products should be inspected visually for particulate matter and discoloration  
10 prior to administration, whenever solution and container permit. If either of these conditions exist,  
11 the vaccine should not be administered.

12 When withdrawing a dose from a stoppered vial, do not remove either the stopper or the metal  
13 seal holding it in place. Use a separate sterile needle and syringe for each injection. Using a sterile  
14 needle and syringe, withdraw the 0.5 mL dose of vaccine from the single-dose vial and administer  
15 the vaccine to the individual. Changing needles between withdrawing the vaccine from the vial  
16 and injecting it into a recipient is not necessary unless the needle has been damaged or  
17 contaminated.

18 Adacel vaccine should not be combined through reconstitution or mixed with any other vaccine.

19 **2.2 Administration, Dose and Schedule**

20 Adacel vaccine is administered as a single 0.5 mL intramuscular injection into the deltoid muscle  
21 of the upper arm.

22 Do not administer this product intravenously, subcutaneously or intradermally.

23 There are no data to support repeat administration of Adacel vaccine.

24 Five years should have elapsed since the recipient's last dose of tetanus toxoid, diphtheria toxoid  
25 and/or pertussis containing vaccine and the administration of Adacel vaccine.

26

27 **2.3 Additional Dosing Information**

28 **Primary series:** The safety and effectiveness of Adacel vaccine used as a primary series or to  
29 complete the primary series, for diphtheria, tetanus, or pertussis has not been demonstrated.

30 **Wound management:** If tetanus prophylaxis is needed for wound management, Adacel may be  
31 given if no previous dose of any Tetanus Toxoid, Reduced Diphtheria Toxoid and Acellular  
32 Pertussis Vaccine, Adsorbed (Tdap) has been administered.

33 **3 DOSAGE FORMS AND STRENGTHS**

34 Adacel vaccine is a suspension for injection (0.5 mL dose) available in 0.5 mL single-dose vials  
35 and prefilled syringes. [See *DOSAGE AND ADMINISTRATION (2.2)* and *HOW*  
36 *SUPPLIED/STORAGE AND HANDLING (16).*]

37 **4 CONTRAINDICATIONS**

38 **4.1 Hypersensitivity**

39 A severe allergic reaction (eg, anaphylaxis) after a previous dose of any tetanus toxoid, diphtheria  
40 toxoid or pertussis containing vaccine or any other component of this vaccine is a contraindication  
41 to administration of Adacel vaccine. [See *DESCRIPTION (11).*] Because of uncertainty as to  
42 which component of the vaccine may be responsible, none of the components should be  
43 administered. Alternatively, such individuals may be referred to an allergist for evaluation if  
44 further immunizations are to be considered.

45 **4.2 Encephalopathy**

46 Encephalopathy (eg, coma, prolonged seizures, or decreased level of consciousness) within 7 days  
47 of a previous dose of a pertussis containing vaccine not attributable to another identifiable cause is  
48 a contraindication to administration of any pertussis containing vaccine, including  
49 Adacel vaccine.

50 **5 WARNINGS AND PRECAUTIONS**

51 **5.1 Management of Acute Allergic Reactions**

52 Epinephrine hydrochloride solution (1:1,000) and other appropriate agents and equipment must be  
53 available for immediate use in case an anaphylactic or acute hypersensitivity reaction occurs.

#### 54 **5.2 Latex**

55 For one presentation of Adacel, the tip caps of the prefilled syringes may contain natural rubber  
56 latex, which may cause allergic reactions in latex sensitive individuals. The vial stopper is not  
57 made with natural rubber latex. [See *HOW SUPPLIED/STORAGE AND HANDLING (16)*.]

#### 58 **5.3 Guillain-Barré Syndrome and Brachial Neuritis**

59 A review by the Institute of Medicine found evidence for acceptance of a causal relation between  
60 tetanus toxoid and both brachial neuritis and Guillain-Barré syndrome. (1) If Guillain-Barré  
61 syndrome occurred within 6 weeks of receipt of prior vaccine containing tetanus toxoid, the  
62 risk for Guillain-Barré syndrome may be increased following a dose of Adacel vaccine.

#### 63 **5.4 Progressive or Unstable Neurologic Disorders**

64 Progressive or unstable neurologic conditions are reasons to defer Adacel. It is not known whether  
65 administration of Adacel to persons with an unstable or progressive neurologic disorder might  
66 hasten manifestations of the disorder or affect the prognosis. Administration of Adacel to persons  
67 with an unstable or progressive neurologic disorder may result in diagnostic confusion between  
68 manifestations of the underlying illness and possible adverse effects of vaccination.

#### 69 **5.5 Arthus-Type Hypersensitivity**

70 Persons who experienced an Arthus-type hypersensitivity reaction following a prior dose of a  
71 tetanus toxoid-containing vaccine should not receive Adacel unless at least 10 years have elapsed  
72 since the last dose of a tetanus toxoid containing vaccine.

#### 73 **5.6 Altered Immunocompetence**

74 If Adacel vaccine is administered to immunocompromised persons, including persons receiving  
75 immunosuppressive therapy, the expected immune response may not be obtained. [See *Drug*  
76 *Interactions (7.2)*.]

77 **5.7 Syncope**

78 Syncope (fainting) can occur in association with administration of injectable vaccine, including  
79 Adacel. Procedures should be in place to prevent falling injury and manage syncopal reactions.

80 **6 ADVERSE REACTIONS**

81 **6.1 Clinical Trials Experience**

82 Because clinical trials are conducted under widely varying conditions, adverse reaction rates  
83 observed in the clinical trials of a vaccine cannot be directly compared to rates in the clinical trials  
84 of another vaccine and may not reflect the rates observed in practice. The adverse reaction  
85 information from clinical trials does, however, provide a basis for identifying the adverse events  
86 that appear to be related to vaccine use and for approximating rates of those events. As with any  
87 vaccine, there is the possibility that broad use of Adacel vaccine could reveal adverse reactions  
88 not observed in clinical trials.

89 The safety of Adacel vaccine was evaluated in 5 clinical studies. A total of 7,143 individuals 10  
90 through 64 years of age inclusive (4,695 adolescents 10 through 17 years of age and, 2,448 adults  
91 18 through 64 years of age) received a single dose of Adacel vaccine.

92 Clinical study Td506 was a randomized, observer-blind, active controlled trial that enrolled  
93 adolescents 11 through 17 years of age (Adacel vaccine N = 1,184; Td vaccine N = 792) and  
94 adults 18 through 64 years of age (Adacel vaccine N = 1,752; Td vaccine N = 573). Study  
95 participants had not received tetanus or diphtheria containing vaccines within the previous 5  
96 years. Solicited local and systemic reactions and unsolicited adverse events were monitored daily  
97 for 14 days post-vaccination using a diary card. From days 14-28 post-vaccination, information on  
98 adverse events necessitating a medical contact, such as a telephone call, visit to an emergency  
99 room, physician's office or hospitalization, was obtained via telephone interview or at an interim  
100 clinic visit. From days 28 to 6 months post-vaccination, participants were monitored for  
101 unexpected visits to a physician's office or to an emergency room, onset of serious illness and  
102 hospitalizations. Information regarding adverse events that occurred in the 6 month post-  
103 vaccination time period was obtained from participants via telephone contact. At least 96% of  
104 participants completed the 6-month follow-up evaluation.

105 **Solicited Adverse Events in the US Adolescent and Adult Study (Td506)**

106 The frequency of selected solicited adverse events (erythema, swelling, pain and fever) occurring  
107 during days 0-14 following vaccination with Adacel vaccine or Td vaccine in adolescents 11  
108 through 17 years of age and adults 18 through 64 years of age are presented in [Table 1](#). Most of  
109 these events were reported at a similar frequency in recipients of both Adacel vaccine and Td  
110 vaccine. Pain at the injection site was the most common adverse reaction in 62.9% to 77.8% of all  
111 vaccinees. In addition, overall rates of pain were higher in adolescent recipients of Adacel vaccine  
112 compared to Td vaccine recipients. Rates of moderate and severe pain in adolescents did not  
113 significantly differ between the Adacel vaccine and Td vaccine groups. Among adults the rates of  
114 pain, after receipt of Adacel vaccine or Td vaccine, did not significantly differ. Fever of 38°C and  
115 higher was uncommon, although in the adolescent age group, it occurred significantly more  
116 frequently in Adacel vaccine recipients than Td vaccine recipients.

117 **Table 1: Frequencies of Solicited Injection Site Reactions and Fever for Adolescents and**  
 118 **Adults, Days 0-14, Following Vaccination With Adacel Vaccine or Td Vaccine in Study**  
 119 **Td506**

Adverse Event*		Adolescents 11-17 years		Adults 18-64 years	
		Adacel N <sup>†</sup> = 1,170-1,175 (%)	Td <sup>‡</sup> N <sup>†</sup> = 783-787 (%)	Adacel N <sup>†</sup> = 1,688-1,698 (%)	Td <sup>‡</sup> N <sup>†</sup> = 551-561 (%)
Injection Site Pain	Any	77.8 <sup>§</sup>	71.0	65.7	62.9
	Moderate**	18.0	15.6	15.1	10.2
	Severe <sup>††</sup>	1.5	0.6	1.1	0.9
Injection Site Swelling	Any	20.9	18.3	21.0	17.3
	Moderate**				
	1.0 to 3.4 cm	6.5	5.7	7.6	5.4
	Severe <sup>††</sup>				
	≥3.5 cm	6.4	5.5	5.8	5.5
	≥5 cm (2 inches)	2.8	3.6	3.2	2.7
Injection Site Erythema	Any	20.8	19.7	24.7	21.6
	Moderate**				
	1.0 to 3.4 cm	5.9	4.6	8.0	8.4
	Severe <sup>††</sup>				
	≥3.5 cm	6.0	5.3	6.2	4.8
	≥5 cm (2 inches)	2.7	2.9	4.0	3.0
Fever	≥38.0°C (≥100.4°F)	5.0 <sup>§</sup>	2.7	1.4	1.1
	≥38.8°C to ≤39.4°C (≥102.0°F to ≤103.0°F)	0.9	0.6	0.4	0.2
	≥39.5°C (≥103.1°F)	0.2	0.1	0.0	0.2

\* The study sample size was designed to detect >10% differences between Adacel and Td vaccines for events of 'Any' intensity.

† N = number of participants with available data.

‡ Tetanus and Diphtheria Toxoids Adsorbed for Adult Use manufactured by Sanofi Pasteur Inc., Swiftwater, PA.

§ Adacel vaccine did not meet the non-inferiority criterion for rates of ‘Any’ Pain in adolescents compared to Td vaccine rates (upper limit of the 95% CI on the difference for Adacel vaccine minus Td vaccine was 10.7% whereas the criterion was <10%). For ‘Any’ Fever the non-inferiority criteria was met, however, ‘Any’ Fever was statistically higher in adolescents receiving Adacel vaccine.

\*\* Interfered with activities, but did not necessitate medical care or absenteeism.

†† Incapacitating, prevented the performance of usual activities, may have/or did necessitate medical care or absenteeism.

120 The frequency of other solicited adverse events (days 0-14) are presented in [Table 2](#). The rates of  
121 these events following Adacel vaccine were comparable with those observed with Td vaccine.

122 Headache was the most frequent systemic reaction and was usually of mild to moderate intensity.



123 **Table 2: Frequencies of Other Solicited Adverse Events for Adolescents and Adults, Days 0-**  
124 **14, Following Vaccination With Adacel Vaccine or Td Vaccine in Study Td506**

Adverse Event		Adolescents 11-17 years		Adults 18-64 years	
		Adacel N* = 1,174-1,175 (%)	Td† N* = 787 (%)	Adacel N* = 1,697-1,698 (%)	Td† N* = 560-561 (%)
Headache	Any	43.7	40.4	33.9	34.1
	Moderate‡	14.2	11.1	11.4	10.5
	Severe§	2.0	1.5	2.8	2.1
Body Ache or Muscle Weakness	Any	30.4	29.9	21.9	18.8
	Moderate‡	8.5	6.9	6.1	5.7
	Severe§	1.3	0.9	1.2	0.9
Tiredness	Any	30.2	27.3	24.3	20.7
	Moderate‡	9.8	7.5	6.9	6.1
	Severe§	1.2	1.0	1.3	0.5
Chills	Any	15.1	12.6	8.1	6.6
	Moderate‡	3.2	2.5	1.3	1.6
	Severe§	0.5	0.1	0.7	0.5
Sore and Swollen Joints	Any	11.3	11.7	9.1	7.0
	Moderate‡	2.6	2.5	2.5	2.1
	Severe§	0.3	0.1	0.5	0.5
Nausea	Any	13.3	12.3	9.2	7.9
	Moderate‡	3.2	3.2	2.5	1.8
	Severe§	1.0	0.6	0.8	0.5
Lymph Node Swelling	Any	6.6	5.3	6.5	4.1
	Moderate‡	1.0	0.5	1.2	0.5
	Severe§	0.1	0.0	0.1	0.0
Diarrhea	Any	10.3	10.2	10.3	11.3
	Moderate‡	1.9	2.0	2.2	2.7
	Severe§	0.3	0.0	0.5	0.5
Vomiting	Any	4.6	2.8	3.0	1.8
	Moderate‡	1.2	1.1	1.0	0.9
	Severe§	0.5	0.3	0.5	0.2
Rash	Any	2.7	2.0	2.0	2.3

\* N = number of participants with available data.

† Tetanus and Diphtheria Toxoids Adsorbed for Adult Use manufactured by Sanofi Pasteur Inc., Swiftwater, PA.

‡ Interfered with activities, but did not necessitate medical care or absenteeism.

§ Incapacitating, prevented the performance of usual activities, may have/or did necessitate medical care or absenteeism.

125 Injection site and systemic solicited reactions occurred at similar rates in Adacel vaccine and  
126 Td vaccine recipients in the 3 day post-vaccination period. Most injection site reactions occurred  
127 within the first 3 days after vaccination (with a mean duration of less than 3 days). The rates of  
128 unsolicited adverse events reported from days 14-28 post-vaccination were comparable between  
129 the two vaccine groups, as were the rates of unsolicited adverse events from day 28 through 6  
130 months. There were no spontaneous reports of extensive limb swelling of the injected limb in  
131 study Td506, nor in the other three studies which also contributed to the safety database for  
132 Adacel vaccine.

### 133 **Injection Site and Systemic Reactions When Given With Hepatitis B Vaccine**

134 In the concomitant vaccination study with Adacel and Hepatitis B vaccines [see *Clinical*  
135 *Studies (14)*], injection site and systemic adverse events were monitored daily for 14 days post-  
136 vaccination using a diary card. Injection site adverse events were only monitored at site/arm of  
137 Adacel vaccine administration. Unsolicited reactions (including immediate reactions, serious  
138 adverse events and events that elicited seeking medical attention) were collected at a clinic visit or  
139 via telephone interview for the duration of the trial, ie, up to 6 months post-vaccination.

140 The rates reported for fever and injection site pain (at the Adacel vaccine administration site) were  
141 similar when Adacel and Hep B vaccines were given concurrently or separately. However, the  
142 rates of injection site erythema (23.4% for concomitant vaccination and 21.4% for separate  
143 administration) and swelling (23.9% for concomitant vaccination and 17.9% for separate  
144 administration) at the Adacel vaccine administration site were increased when co-administered.  
145 Swollen and/or sore joints were reported by 22.5% for concomitant vaccination and 17.9% for  
146 separate administration. The rates of generalized body aches in the individuals who reported  
147 swollen and/or sore joints were 86.7% for concomitant vaccination and 72.2% for separate  
148 administration. Most joint complaints were mild in intensity with a mean duration of 1.8 days.  
149 The incidence of other solicited and unsolicited adverse events were not different between the  
150 2 study groups.

### 151 **Injection Site and Systemic Reactions When Given With Trivalent Inactivated Influenza** 152 **Vaccine (TIV)**

153 In the concomitant vaccination study with Adacel vaccine and trivalent inactivated influenza  
154 vaccine [see *Clinical Studies (14)*], injection site and systemic adverse events were monitored for

155 14 days post-vaccination using a diary card. All unsolicited reactions occurring through day 14  
156 were collected. From day 14 to the end of the trial, ie, up to 84 days, only events that elicited  
157 seeking medical attention were collected.

158 The rates of fever and injection site erythema and swelling were similar for recipients of  
159 concurrent and separate administration of Adacel vaccine and TIV. However, pain at the Adacel  
160 vaccine injection site occurred at statistically higher rates following concurrent administration  
161 (66.6%) versus separate administration (60.8%). The rates of sore and/or swollen joints were  
162 13% for concurrent administration and 9% for separate administration. Most joint complaints  
163 were mild in intensity with a mean duration of 2.0 days. The incidence of other solicited and  
164 unsolicited adverse events were similar between the 2 study groups.

### 165 **Additional Studies**

166 In an additional study, 1,806 adolescents 11 through 17 years of age received Adacel vaccine as  
167 part of the lot consistency study used to support Adacel vaccine licensure. This study was a  
168 randomized, double-blind, multi-center trial designed to assess lot consistency as measured by the  
169 safety and immunogenicity of 3 lots of Adacel vaccine when given as a booster dose to  
170 adolescents 11 through 17 years of age inclusive. Local and systemic adverse events were  
171 monitored for 14 days post-vaccination using a diary card. Unsolicited adverse events and serious  
172 adverse events were collected for 28 days post-vaccination. Pain was the most frequently reported  
173 local adverse event occurring in approximately 80% of all participants. Headache was the most  
174 frequently reported systemic event occurring in approximately 44% of all participants. Sore  
175 and/or swollen joints were reported by approximately 14% of participants. Most joint complaints  
176 were mild in intensity with a mean duration of 2.0 days.

177 An additional 962 adolescents and adults received Adacel vaccine in three supportive Canadian  
178 studies used as the basis for licensure in other countries. Within these clinical trials, the rates of  
179 local and systemic reactions following Adacel vaccine were similar to those reported in the four  
180 principal trials in the US with the exception of a higher rate (86%) of adults experiencing ‘any’  
181 local injection site pain. The rate of severe pain (0.8%), however, was comparable to the rates  
182 reported in four principal trials conducted in the US. There was one spontaneous report of whole-  
183 arm swelling of the injected limb among the 277 Td vaccine recipients, and two spontaneous  
184 reports among the 962 Adacel vaccine recipients in the supportive Canadian studies.

185 An additional study, Td519, enrolled 1,302 individuals in an open label, two-arm, multi-center  
186 trial (651 subjects in each group) to evaluate the safety and immunogenicity of a single dose of  
187 Adacel administered to persons 10 to <11 years of age compared to persons 11 to <12 years of  
188 age. Immediate reactions were monitored for 20 minutes post-vaccination. Solicited local and  
189 systemic adverse events were monitored for 7 days post-vaccination using a diary card.  
190 Unsolicited and serious adverse events were collected for approximately 30 days post-  
191 vaccination. Similar rates of immediate, solicited and unsolicited adverse reactions were reported  
192 in each of the two age cohorts. One serious adverse event, not related to vaccination, was reported  
193 in the younger age group.

#### 194 **Serious Adverse Events in All Safety Studies**

195 In all the studies, participants were monitored for serious adverse events throughout the duration  
196 of the study.  
197 Throughout the 6-month follow-up period in study Td506, serious adverse events were reported in  
198 1.5% of Adacel vaccine recipients and in 1.4% of Td vaccine recipients. Two serious adverse  
199 events in adults were neuropathic events that occurred within 28 days of Adacel vaccine  
200 administration; one severe migraine with unilateral facial paralysis and one diagnosis of nerve  
201 compression in neck and left arm. Similar or lower rates of serious adverse events were reported  
202 in the other trials in participants up to 64 years of age and no additional neuropathic events were  
203 reported.

## 204 **6.2 Postmarketing Experience**

205 The following adverse events of Adacel have been spontaneously reported in the US and other  
206 countries. Because these events are reported voluntarily from a population of uncertain size, it  
207 may not be possible to reliably estimate their frequency or establish a causal relationship to  
208 vaccine exposure.

209 The following adverse events were included based on one or more of the following factors:  
210 severity, frequency of reporting or strength of evidence for a causal relationship to Adacel  
211 vaccine.

#### 212 ***Immune system disorders***

213 Anaphylactic reaction, hypersensitivity reaction (angioedema, edema, rash, hypotension)

214 ☒ ***Nervous system disorders***

215 Paraesthesia, hypoesthesia, Guillain-Barré syndrome, brachial neuritis, facial palsy,  
216 convulsion, syncope, myelitis

217 ☒ ***Cardiac disorders***

218 Myocarditis

219 ☒ ***Skin and subcutaneous tissue disorders***

220 Pruritus, urticaria

221 ☒ ***Musculoskeletal and connective tissue disorders***

222 Myositis, muscle spasm

223 ☒ ***General disorders and administration site conditions***

224 Large injection site reactions (>50 mm), extensive limb swelling from the injection site  
225 beyond one or both joints  
226 Injection site bruising, sterile abscess

227 **7 DRUG INTERACTIONS**

228 **7.1 Concomitant Vaccine Administration**

229 When Adacel vaccine is administered concomitantly with other injectable vaccines or Tetanus  
230 Immune Globulin, they should be given with separate syringes and at different injection sites.  
231 Adacel should not be mixed with any other vaccine in the same syringe or vial.

232 In clinical studies, Adacel vaccine was administered concomitantly with one of the following US-  
233 licensed vaccines: Hepatitis B (10 mcg, two dose regimen) or trivalent inactivated influenza  
234 vaccines (TIV). [See *Adverse Reactions (6.1)* and *Clinical Studies (14)*.]

235 **Hepatitis B Vaccine**

236 Concomitant immunization of Adacel vaccine with Hepatitis B vaccine did not result in reduced  
237 antibody responses to any of the antigens from either vaccine.

238           **Trivalent Inactivated Influenza Vaccine (TIV)**

239   No interference in tetanus and diphtheria seroprotection rates and responses to influenza vaccine,  
240   detoxified pertussis toxin (PT), fimbriae types 2 and 3 (FIM) or filamentous hemagglutinin (FHA)  
241   were observed when Adacel vaccine was administered concomitantly with TIV compared to  
242   separate administration. A lower pertactin (PRN) GMC was observed when Adacel vaccine was  
243   administered concomitantly with TIV compared to separate administration.

244   **7.2 Immunosuppressive Treatments**

245   Immunosuppressive therapies, including irradiation, antimetabolites, alkylating agents, cytotoxic  
246   drugs and corticosteroids (used in greater than physiologic doses), may reduce the immune  
247   response to vaccines. [See *Warnings and Precautions* (5.6).]

248

## 249 **8 USE IN SPECIFIC POPULATIONS**

### 250 **8.1 Pregnancy**

#### 251 **Pregnancy Category C**

252 Animal reproduction studies have not been conducted with Adacel vaccine. It is also not known  
253 whether Adacel vaccine can cause fetal harm when administered to a pregnant woman or can  
254 affect reproduction capacity. Adacel vaccine should be given to a pregnant woman only if clearly  
255 needed.

256 Animal fertility studies have not been conducted with Adacel vaccine. The effect of Adacel  
257 vaccine on embryo-fetal and pre-weaning development was evaluated in two developmental  
258 toxicity studies using pregnant rabbits. Animals were administered Adacel vaccine twice prior to  
259 gestation, during the period of organogenesis (gestation day 6) and later during pregnancy on  
260 gestation day 29, 0.5 mL/rabbit/occasion (a 17-fold increase compared to the human dose of  
261 Adacel vaccine on a body weight basis), by intramuscular injection. No adverse effects on  
262 pregnancy, parturition, lactation, embryo-fetal or pre-weaning development were observed. There  
263 were no vaccine related fetal malformations or other evidence of teratogenesis noted in this study.

#### 264 **Registry of Receipt of Adacel Vaccine During Pregnancy**

265 Sanofi Pasteur Inc. maintains a surveillance registry to collect data on pregnancy outcomes and  
266 newborn health status outcomes following vaccination with Adacel vaccine during pregnancy.  
267 Women who receive Adacel vaccine during pregnancy are encouraged to contact directly or have  
268 their health-care professional contact Sanofi Pasteur Inc. at 1-800-822-2463 (1-800-VACCINE).

### 269 **8.3 Nursing Mothers**

270 It is not known whether Adacel vaccine is excreted in human milk. Because many drugs are  
271 excreted in human milk, caution should be exercised when Adacel vaccine is given to a nursing  
272 woman.

273

**274 8.4 Pediatric Use**

275 Adacel vaccine is not approved for individuals less than 10 years of age. Safety and effectiveness  
276 of Adacel vaccine in persons less than 10 years of age have not been established.

**277 8.5 Geriatric Use**

278 Adacel vaccine is not approved for use in individuals 65 years of age and older.  
279 In a clinical study, individuals 65 years of age and older received a single dose of Adacel vaccine.  
280 Based on pre-specified criteria, persons 65 years of age and older who received a dose of Adacel  
281 vaccine had lower geometric mean concentrations of antibodies to PT, PRN and FIM when  
282 compared to infants who had received a primary series of DAPTACEL<sup>®</sup>, Diphtheria and Tetanus  
283 Toxoids and Acellular Pertussis Vaccine Adsorbed (DTaP). [See Section 14 for description of  
284 DAPTACEL vaccine.]

**285 11 DESCRIPTION**

286 Adacel vaccine is a sterile isotonic suspension of tetanus and diphtheria toxoids and pertussis  
287 antigens adsorbed on aluminum phosphate, for intramuscular injection.  
288 Each 0.5 mL dose contains 5 Lf tetanus toxoid (T), 2 Lf diphtheria toxoid (d), and acellular  
289 pertussis antigens [2.5 mcg detoxified pertussis toxin (PT), 5 mcg filamentous hemagglutinin  
290 (FHA), 3 mcg pertactin (PRN), 5 mcg fimbriae types 2 and 3 (FIM)]. Other ingredients per 0.5  
291 mL dose include 1.5 mg aluminum phosphate (0.33 mg aluminum) as the adjuvant, ≤5 mcg  
292 residual formaldehyde, <50 ng residual glutaraldehyde and 3.3 mg (0.6% v/v) 2-phenoxyethanol  
293 (not as a preservative). The antigens are the same as those in DAPTACEL vaccine; however,  
294 Adacel vaccine is formulated with reduced quantities of diphtheria and detoxified PT.  
295 The acellular pertussis vaccine components are produced from *Bordetella pertussis* cultures  
296 grown in Stainer-Scholte medium (2) modified by the addition of casamino acids and dimethyl-  
297 beta-cyclodextrin. PT, FHA and PRN are isolated separately from the supernatant culture  
298 medium. FIM are extracted and co-purified from the bacterial cells. The pertussis antigens are  
299 purified by sequential filtration, salt-precipitation, ultrafiltration and chromatography. PT is  
300 detoxified with glutaraldehyde, FHA is treated with formaldehyde, and the residual aldehydes are  
301 removed by ultrafiltration. The individual antigens are adsorbed onto aluminum phosphate.  
302 The tetanus toxin is produced from *Clostridium tetani* grown in modified Mueller-Miller



303 casamino acid medium without beef heart infusion. (3) Tetanus toxin is detoxified with  
304 formaldehyde and purified by ammonium sulfate fractionation and diafiltration. *Corynebacterium*  
305 *diphtheriae* is grown in modified Mueller's growth medium. (4) After purification by ammonium  
306 sulfate fractionation, diphtheria toxin is detoxified with formaldehyde and diafiltered.  
307 The adsorbed diphtheria, tetanus and acellular pertussis components are combined with aluminum  
308 phosphate (as adjuvant), 2-phenoxyethanol (not as a preservative) and water for injection. Adacel  
309 vaccine does not contain a preservative.  
310 In the guinea pig potency test, the tetanus component induces at least 2 neutralizing units/mL of  
311 serum and the diphtheria component induces at least 0.5 neutralizing units/mL of serum. The  
312 potency of the acellular pertussis vaccine components is evaluated by the antibody response of  
313 immunized mice to detoxified PT, FHA, PRN and FIM as measured by enzyme-linked  
314 immunosorbent assay (ELISA).  
315 Diphtheria and tetanus toxoids are individually adsorbed onto aluminum phosphate.  
316

317 **12 CLINICAL PHARMACOLOGY**

318 **12.1 Mechanism of Action**

319 **Tetanus**

320 Tetanus is a disease manifested primarily by neuromuscular dysfunction caused by a potent  
321 exotoxin released by *C tetani*.

322 Protection against disease is due to the development of neutralizing antibodies to tetanus toxin. A  
323 serum tetanus antitoxin level of at least 0.01 IU/mL, measured by neutralization assay is  
324 considered the minimum protective level. (5) (6)

325 **Diphtheria**

326 Diphtheria is an acute toxin-mediated disease caused by toxigenic strains of *C diphtheriae*.

327 Protection against disease is due to the development of neutralizing antibodies to diphtheria toxin.  
328 A serum diphtheria antitoxin level of 0.01 IU/mL is the lowest level giving some degree of  
329 protection. Antitoxin levels of at least 0.1 IU/mL are generally regarded as protective. (5) Levels  
330 of 1.0 IU/mL have been associated with long-term protection. (7)

331 **Pertussis**

332 Pertussis (whooping cough) is a respiratory disease caused by *B pertussis*. This Gram-negative  
333 coccobacillus produces a variety of biologically active components, though their role in either the  
334 pathogenesis of, or immunity to, pertussis has not been clearly defined.

335 **13 NON-CLINICAL TOXICOLOGY**

336 **13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility**

337 Adacel vaccine has not been evaluated for carcinogenic or mutagenic potential, or impairment of  
338 fertility.

339

## 340 **14 CLINICAL STUDIES**

341 The efficacy of the tetanus toxoid and diphtheria toxoid used in Adacel vaccine was based on the  
342 immune response to these antigens compared to a US licensed Tetanus and Diphtheria Toxoids  
343 Adsorbed For Adult Use (Td) vaccine manufactured by Sanofi Pasteur Inc., Swiftwater, PA. The  
344 primary measures for immune response to the diphtheria and tetanus toxoids were the percentage  
345 of participants attaining an antibody level of at least 0.1 IU/mL.

346 The efficacy of the pertussis antigens used in Adacel vaccine was inferred based on a comparison  
347 of pertussis antibody levels achieved in recipients of a single booster dose of Adacel vaccine with  
348 those obtained in infants after three doses of DAPTACEL vaccine. In the Sweden I Efficacy Trial,  
349 three doses of DAPTACEL vaccine were shown to confer a protective efficacy of 84.9% (95%  
350 CI: 80.1%, 88.6%) against WHO defined pertussis (21 days of paroxysmal cough with laboratory-  
351 confirmed *B pertussis* infection or epidemiological link to a confirmed case). The protective  
352 efficacy against mild pertussis (defined as at least one day of cough with laboratory-confirmed  
353 *B pertussis* infection) was 77.9% (95% CI: 72.6%, 82.2%). (8)

354 In addition, the ability of Adacel vaccine to elicit a booster response (defined as rise in antibody  
355 concentration after vaccination) to the tetanus, diphtheria and pertussis antigens following  
356 vaccination was evaluated. The demonstration of a booster response depended on the antibody  
357 concentration to each antigen as established based on the 95<sup>th</sup> percentile of the pre-vaccination  
358 antibody concentrations observed in historical clinical trials with Adacel vaccine.

### 359 **14.1 Immunological Evaluation in Adolescents and Adults, 10 Through 64 Years of** 360 **Age**

361 Study Td506 was a comparative, multi-center, randomized, observer-blind, controlled trial which  
362 enrolled 4,480 participants; 2,053 adolescents (11 through 17 years of age) and 2,427 adults (18  
363 through 64 years of age). Enrollment was stratified by age to ensure adequate representation  
364 across the entire age range. Participants had not received a tetanus or diphtheria toxoid containing  
365 vaccine within the previous 5 years. After enrollment participants were randomized to receive one  
366 dose of either Adacel vaccine or Td vaccine. A total of 4,461 randomized participants were  
367 vaccinated. The per-protocol immunogenicity subset included 1,270 Adacel vaccine recipients  
368 and 1,026 Td vaccine recipients. Sera were obtained before and approximately 35 days after

369 vaccination. [Blinding procedures for safety assessments are described in *ADVERSE REACTIONS*  
370 [\(6\)](#).]

371 Demographic characteristics were similar within age groups and between the vaccine groups. A  
372 total of 76% of the adolescents and 1.1% of the adults reported a history of receiving 5 previous  
373 doses of diphtheria-tetanus-pertussis containing vaccines. Anti-tetanus and anti-diphtheria  
374 seroprotection rates ( $\geq 0.1$  IU/mL) and booster response rates were comparable between Adacel  
375 and Td vaccines. (See [Table 3](#) and [Table 4](#).) Adacel vaccine induced pertussis antibody levels that  
376 were non-inferior to those of Swedish infants who received three doses of DAPTACEL vaccine.  
377 (See [Table 5](#).) Acceptable booster responses to each of the pertussis antigens were also  
378 demonstrated, ie, the percentage of participants with a booster response exceeded the pre-defined  
379 lower limit. (See [Table 6](#).)

380 **Table 3: Pre-vaccination and Post-vaccination Antibody Responses and Booster Response**  
 381 **Rates to Tetanus Toxoid Following Adacel Vaccine as Compared to Td Vaccine in**  
 382 **Adolescents and Adults 11 Through 64 Years of Age**

			Tetanus Antitoxin (IU/mL)				
			Pre-vaccination		1 Month Post-vaccination		
Age Group (years)	Vaccine	N*	% ≥0.10 (95% CI)	% ≥1.0 (95% CI)	% ≥0.10 (95% CI)	% ≥1.0 (95% CI)	% Booster† (95% CI)
11-17	Adacel	527	99.6 (98.6, 100.0)	44.6 (40.3, 49.0)	100.0‡ (99.3, 100.0)	99.6§ (98.6, 100.0)	91.7‡ (89.0, 93.9)
	Td**	516	99.2 (98.0, 99.8)	43.8 (39.5, 48.2)	100.0 (99.3, 100.0)	99.4 (98.3, 99.9)	91.3 (88.5, 93.6)
18-64	Adacel	742-743	97.3 (95.9, 98.3)	72.9 (69.6, 76.1)	100.0‡ (99.5, 100.0)	97.8§ (96.5, 98.8)	63.1‡ (59.5, 66.6)
	Td**	509	95.9 (93.8, 97.4)	70.3 (66.2, 74.3)	99.8 (98.9, 100.0)	98.2 (96.7, 99.2)	66.8 (62.5, 70.9)

\* N = number of participants in the per-protocol population with available data.

† Booster response is defined as: A four-fold rise in antibody concentration, if the pre-vaccination concentration was equal to or below the cut-off value and a two-fold rise in antibody concentration if the pre-vaccination concentration was above the cut-off value. The cut-off value for tetanus was 2.7 IU/mL.

‡ Seroprotection rates at ≥0.10 IU/mL and booster response rates to Adacel vaccine were non-inferior to Td vaccine (upper limit of the 95% CI on the difference for Td vaccine minus Adacel vaccine <10%).

§ Seroprotection rates at ≥1.0 IU/mL were not prospectively defined as a primary endpoint.

\*\* Tetanus and Diphtheria Toxoids Adsorbed for Adult Use manufactured by Sanofi Pasteur Inc., Swiftwater, PA.

383 **Table 4: Pre-vaccination and Post-vaccination Antibody Responses and Booster Response**  
 384 **Rates to Diphtheria Toxoid Following Adacel Vaccine as Compared to Td Vaccine in**  
 385 **Adolescents and Adults 11 Through 64 Years of Age**

			Diphtheria Antitoxin (IU/mL)				
			Pre-vaccination		1 Month Post-vaccination		
Age Group (years)	Vaccine	N*	% ≥0.10 (95% CI)	% ≥1.0 (95% CI)	% ≥0.10 (95% CI)	% ≥1.0 (95% CI)	% Booster <sup>†</sup> (95% CI)
11-17	Adacel	527	72.5 (68.5, 76.3)	15.7 (12.7, 19.1)	99.8 <sup>‡</sup> (98.9, 100.0)	98.7 <sup>§</sup> (97.3, 99.5)	95.1 <sup>‡</sup> (92.9, 96.8)
	Td**	515-516	70.7 (66.5, 74.6)	17.3 (14.1, 20.8)	99.8 (98.9, 100.0)	98.4 (97.0, 99.3)	95.0 (92.7, 96.7)
18-64	Adacel	739-741	62.6 (59.0, 66.1)	14.3 (11.9, 17.0)	94.1 <sup>‡</sup> (92.1, 95.7)	78.0 <sup>§</sup> (74.8, 80.9)	87.4 <sup>‡</sup> (84.8, 89.7)
	Td**	506-507	63.3 (59.0, 67.5)	16.0 (12.9, 19.5)	95.1 (92.8, 96.8)	79.9 (76.1, 83.3)	83.4 (79.9, 86.5)

\* N = number of participants in the per-protocol population with available data.

† Booster response is defined as: A four-fold rise in antibody concentration, if the pre-vaccination concentration was equal to or below the cut-off value and a two-fold rise in antibody concentration if the pre-vaccination concentration was above the cut-off value. The cut-off value for diphtheria was 2.56 IU/mL.

‡ Seroprotection rates at ≥0.10 IU/mL and booster response rates to Adacel vaccine were non-inferior to Td vaccine (upper limit of the 95% CI on the difference for Td vaccine minus Adacel vaccine <10%).

§ Seroprotection rates at ≥1.0 IU/mL were not prospectively defined as a primary endpoint.

\*\* Tetanus and Diphtheria Toxoids Adsorbed for Adult Use manufactured by Sanofi Pasteur Inc., Swiftwater, PA.

386 **Table 5: Ratio of Pertussis Antibody Geometric Mean Concentrations (GMCs)<sup>¥</sup> Observed**  
 387 **One Month After a Dose of Adacel Vaccine in Adolescents and Adults 11 Through 64 Years**  
 388 **of Age Compared With Those Observed in Infants One Month Following Vaccination at 2, 4**  
 389 **and 6 Months of Age in the Efficacy Trial With DAPTACEL Vaccine**

	Adolescents 11-17 Years of Age	Adults 18-64 Years of Age
	Adacel*/DAPTACEL <sup>†</sup> GMC Ratio (95% CIs)	Adacel <sup>‡</sup> /DAPTACEL <sup>†</sup> GMC Ratio (95% CIs)
<b>Anti-PT</b>	3.6 (2.8, 4.5) <sup>§</sup>	2.1 (1.6, 2.7) <sup>§</sup>
<b>Anti-FHA</b>	5.4 (4.5, 6.5) <sup>§</sup>	4.8 (3.9, 5.9) <sup>§</sup>
<b>Anti-PRN</b>	3.2 (2.5, 4.1) <sup>§</sup>	3.2 (2.3, 4.4) <sup>§</sup>
<b>Anti-FIM</b>	5.3 (3.9, 7.1) <sup>§</sup>	2.5 (1.8, 3.5) <sup>§</sup>

¥ Antibody GMCs, measured in arbitrary ELISA units were calculated separately for infants, adolescents and adults.

\* N = 524 to 526, number of adolescents in the per-protocol population with available data for Adacel vaccine.

† N = 80, number of infants who received DAPTACEL vaccine with available data post-dose 3 (Sweden Efficacy I).

‡ N = 741, number of adults in the per-protocol population with available data for Adacel vaccine.

§ GMC following Adacel vaccine was non-inferior to GMC following DAPTACEL vaccine (lower limit of 95% CI on the ratio of GMC for Adacel vaccine divided by DAPTACEL vaccine >0.67).

390 **Table 6: Booster Response Rates to the Pertussis Antigens Observed One Month After a**  
391 **Dose of Adacel Vaccine in Adolescents and Adults 11 Through 64 Years of Age**

	Adolescents 11-17 Years of Age		Adults 18-64 Years of Age		Pre-defined Acceptable Rates* %†
	N‡	% (95% CI)	N‡	% (95% CI)	
<b>Anti-PT</b>	524	92.0 (89.3, 94.2)	739	84.4 (81.6, 87.0)	81.2
<b>Anti-FHA</b>	526	85.6 (82.3, 88.4)	739	82.7 (79.8, 85.3)	77.6
<b>Anti-PRN</b>	525	94.5 (92.2, 96.3)	739	93.8 (91.8, 95.4)	86.4
<b>Anti-FIM</b>	526	94.9 (92.6, 96.6)	739	85.9 (83.2, 88.4)	82.4

\* The acceptable response rate for each antigen was defined as the lower limit of the 95% CI for the rate being no more than 10% lower than the response rate observed in previous clinical trials.

† A booster response for each antigen was defined as a four-fold rise in antibody concentration if the pre-vaccination concentration was equal to or below the cut-off value and a two-fold rise in antibody concentration if the pre-vaccination concentration was above the cut-off value. The cut-off values for pertussis antigens were established based on antibody data from both adolescents and adults in previous clinical trials.  
The cut-off values were 85 EU/mL for PT, 170 EU/mL for FHA, 115 EU/mL for PRN and 285 EU/mL for FIM.

‡ N = number of participants in the per-protocol population with available data.



392 Study Td519 assessed the comparative immunogenicity of Adacel administered to adolescents  
393 (10 to <11 years of age and 11 to <12 years of age) [see *Adverse Reactions* (6.1).] In this study  
394 non-inferiority was demonstrated for booster responses to tetanus and diphtheria toxoids, GMCs  
395 to the pertussis antigens (PT, FHA, PRN and FIM) and booster responses to the pertussis antigens  
396 PT, FHA and PRN. For FIM, non-inferiority was not demonstrated as the lower bound of the 95%  
397 CI of the difference in booster response rates (-5.96%) did not meet the predefined criterion (>-  
398 5% when the booster response in the older age group was >95%).

#### 399 **14.2 Concomitant Hepatitis B Vaccine Administration**

400 The concomitant use of Adacel vaccine and hepatitis B (Hep B) vaccine (Recombivax HB®, 10  
401 mcg per dose using a two-dose regimen, manufactured by Merck and Co., Inc) was evaluated in a  
402 multi-center, open-labeled, randomized, controlled study that enrolled 410 adolescents, 11  
403 through 14 years of age inclusive. One group received Adacel and Hep B vaccines concurrently  
404 (N = 206). The other group (N = 204) received Adacel vaccine at the first visit, then 4-6 weeks  
405 later received Hep B vaccine. The second dose of Hep B vaccine was given 4-6 weeks after the  
406 first dose. Serum samples were obtained prior to and 4-6 weeks after Adacel vaccine  
407 administration, as well as 4-6 weeks after the 2<sup>nd</sup> dose of Hep B for all participants. No  
408 interference was observed in the immune responses to any of the vaccine antigens when Adacel  
409 and Hep B vaccines were given concurrently or separately. [See *ADVERSE REACTIONS* (6.1).]

#### 410 **14.3 Concomitant Influenza Vaccine Administration**

411 The concomitant use of Adacel vaccine and trivalent inactivated influenza vaccine (TIV,  
412 Fluzone®, manufactured by Sanofi Pasteur Inc., Swiftwater, PA) was evaluated in a multi-center,  
413 open-labeled, randomized, controlled study conducted in 720 adults, 19-64 years of age inclusive.  
414 In one group, participants received Adacel and TIV vaccines concurrently (N = 359). The other  
415 group received TIV at the first visit, then 4-6 weeks later received Adacel vaccine (N = 361). Sera  
416 were obtained prior to and 4-6 weeks after Adacel vaccine, as well as 4-6 weeks after the TIV.  
417 The immune responses were comparable for concurrent and separate administration of Adacel and  
418 TIV vaccines for diphtheria (percent of participants with seroprotective concentration  $\geq 0.10$   
419 IU/mL and booster responses), tetanus (percent of participants with seroprotective concentration  
420  $\geq 0.10$  IU/mL), pertussis antigens (booster responses and GMCs except lower PRN GMC in the  
421 concomitant group, lower bound of the 90% CI was 0.61 and the pre-specified criterion was

422  $\geq 0.67$ ) and influenza antigens (percent of participants with hemagglutination-inhibition [HI]  
423 antibody titer  $\geq 1:40$  IU/mL and  $\geq 4$ -fold rise in HI titer). Although tetanus booster response rates  
424 were significantly lower in the group receiving the vaccines concurrently versus separately,  
425 greater than 98% of participants in both groups achieved seroprotective levels of  $\geq 0.1$  IU/mL.  
426 [See *ADVERSE REACTIONS (6.1)*.]

427

428 **15 REFERENCES**

429

- 430 1 Stratton KR, et al, editors. Adverse events associated with childhood vaccines; evidence  
431 bearing on causality. Washington: National Academy Press; 1994. p. 67-117.
- 432 2 Stainer DW, et al. A simple chemically defined medium for the production of phase I  
433 Bordetella pertussis. J Gen Microbiol 1970;63:211-20.
- 434 3 Mueller JH, et al. Variable factors influencing the production of tetanus toxin. J Bacteriol  
435 1954;67(3):271-7.
- 436 4 Stainer DW. Production of diphtheria toxin. In: Manclark CR, editor. Proceedings of an  
437 informal consultation on the World Health Organization requirements for diphtheria,  
438 tetanus, pertussis and combined vaccines. United States Public Health Service, Bethesda,  
439 MD. DHHS 91-1174. 1991. p. 7-11.
- 440 5 FDA. Department of Health and Human Services (DHHS). Biological products bacterial  
441 vaccines and toxoids; implementation of efficacy review; proposed rule. Fed Reg  
442 1985;50(240):51002-117.
- 443 6 Wassilak SGF, et al. Tetanus toxoid. In: Plotkin SA, Orenstein WA, Offit PA, editors.  
444 Vaccines. 5th ed. Philadelphia, PA: W.B. Saunders Company; 2008. p. 805-39.
- 445 7 Vitek CR and Wharton M. Diphtheria toxoid. In: Plotkin SA, Orenstein WA, Offit PA,  
446 editors. Vaccines. 5th ed. Philadelphia, PA: W.B. Saunders Company; 2008. p. 139-56.
- 447 8 Gustafsson L, et al. A controlled trial of a two-component acellular, a five-component  
448 acellular and a whole-cell pertussis vaccine. N Engl J Med 1996;334(6):349-55.

449

450

451 **16 HOW SUPPLIED/STORAGE AND HANDLING**

452 Syringe, without needle, 1 dose - NDC No. 49281-400-89 (not made with natural rubber latex); in  
453 package of 5 syringes, NDC No. 49281-400-20.

454 Syringe, without needle, 1 dose - NDC No. 49281-400-88; in package of 5 syringes, NDC No.  
455 49281-400-15. The tip caps of the prefilled syringes may contain natural rubber latex. No other  
456 components are made with natural rubber latex.

457 Vial, 1 dose - NDC No. 49281-400-58; in package of 5 vials; NDC No. 49281-400-05. The vial  
458 stopper is not made with natural rubber latex.

459 Vial, 1 dose - NDC No. 49281-400-58; in package of 10 vials; NDC No. 49281-400-10. The vial  
460 stopper is not made with natural rubber latex.

461 Adacel vaccine should be stored at 2° to 8°C (35° to 46°F). DO NOT FREEZE. Product which  
462 has been exposed to freezing should not be used. Do not use after expiration date shown on the  
463 label.

464 **17 PATIENT COUNSELING INFORMATION**

465 Before administration of Adacel vaccine, health-care providers should inform the patient, parent  
466 or guardian of the benefits and risks of the vaccine and the importance of receiving recommended  
467 booster dose unless a contraindication to further immunization exists.

468 The health-care provider should inform the patient, parent or guardian about the potential for  
469 adverse reactions that have been temporally associated with Adacel vaccine or other vaccines  
470 containing similar components. The health-care provider should provide the Vaccine Information  
471 Statements (VISs) that are required by the National Childhood Vaccine Injury Act of 1986 to be  
472 given with each immunization. The patient, parent or guardian should be instructed to report any  
473 serious adverse reactions to their health-care provider.

474 **Pregnancy Exposure Registry** [See *USE IN SPECIFIC POPULATIONS (8.1).*]

475

476

Printed in XXX

477 Manufactured by:

478 **Sanofi Pasteur Limited**

479 Toronto Ontario Canada

480

481 Distributed by:

482 **Sanofi Pasteur Inc.**

483 Swiftwater PA 18370 USA

484

485 Adacel<sup>®</sup> is a registered trademark of Sanofi, its affiliates and its subsidiaries.

486

R10-XX17 USA

487 **SANOFI PASTEUR** 

488